

NEW SOLUTIONS, Vol. 23(1) 55-83, 2013

Scientific Solutions

**INVESTIGATING LINKS BETWEEN SHALE GAS
DEVELOPMENT AND HEALTH IMPACTS THROUGH A
COMMUNITY SURVEY PROJECT IN PENNSYLVANIA**

**NADIA STEINZOR
WILMA SUBRA
LISA SUMI**

ABSTRACT

Across the United States, the race for new energy sources is picking up speed and reaching more places, with natural gas in the lead. While the toxic and polluting qualities of substances used and produced in shale gas development and the general health effects of exposure are well established, scientific evidence of causal links has been limited, creating an urgent need to understand health impacts. Self-reported survey research documenting the symptoms experienced by people living in proximity to gas facilities, coupled with environmental testing, can elucidate plausible links that warrant both response and further investigation. This method, recently applied to the gas development areas of Pennsylvania, indicates the need for a range of policy and research efforts to safeguard public health.

Keywords: health surveys, shale gas, toxic exposure, hydraulic fracturing, fracking

Public health was not brought into discussions about shale gas extraction at earlier stages; in consequence, the health system finds itself lacking critical information about environmental and public health impacts of the technologies and unable to address concerns by regulators at the federal and state levels, communities, and workers. . . .

—Institute of Medicine at the National Academies of Science [1]

For many years, extracting natural gas from deep shale formations across the United States (such as the Marcellus Shale in the East or the Barnett Shale in Texas) was considered economically and technologically infeasible. More recently, changes in hydraulic fracturing technology and its combination with horizontal drilling have made it possible to drill much deeper and further. Bolstered by declining global oil resources and a strong political push to expand domestic energy production, this has resulted in a boom in shale gas production nationwide and projections of tens or even hundreds of thousands of wells being drilled in the coming decades.

By mid-2012, there were nearly 490,000 producing natural gas wells in the United States, 60,000 more than in 2005 [2]. In Pennsylvania alone, more than 5,900 unconventional oil and gas wells had been drilled, and more than 11,700 had been permitted, between 2005 and September 2012; the pace of expansion has been rapid, with 75 percent of all unconventional wells drilled just in the last two years [3]. The rapid pace of industry expansion is increasingly divergent from the slower pace of scientific understanding of its impacts, as well as policy and regulatory measures to prevent them—in turn raising many questions that have yet to be answered [4]. Further, the limited availability of information has both contributed to public perception and supported industry assertions that health impacts related to oil and gas development are isolated and rare.

Modern-day industrial gas and oil development has many stages, uses a complex of chemicals, and produces large volumes of both wastewater and solid waste, which create the potential for numerous pathways of exposure to substances harmful to health, in particular to air and water pollution [5]. Many reports of negative health impacts by people living in proximity to wells and oil and gas facilities have been documented in the media and through research by organizations [6-8]. In addition, several self-reporting health survey and environmental testing projects have been conducted in response to complaints following pollution events or the establishment of facilities [9-12].

Such short-term projects have been initiated in a research context in which longer-term investigations—particularly ones that seek to establish causal links between health problems and oil and gas development—have historically been narrow and inconsistent [13]. Reflecting growing concern over the need to deepen knowledge among scientists, public agency representatives, and environmental and health professionals, four conferences on the links between shale gas development and human health were convened in just a one-year period (November 2011–November 2012), including those convened by the Graduate School of Public Health at the University of Pittsburgh; by Physicians, Scientists, and Engineers for Healthy Energy; and by the Institute of Medicine of the National Academy of Sciences.

In-depth research on the health impacts of oil and gas development has also begun to appear in the literature. In 2011, a review of more than 600 known chemicals used in natural gas operations concluded that many could cause cancer

and mutations and have long-term health impacts (including on the skin, eyes, and kidneys and on the respiratory, gastrointestinal, brain/nervous, immune, endocrine, and cardiovascular systems) [14]. In early 2012, a study by researchers at the University of Colorado concluded that the toxicity of air emissions near natural gas sites puts residents living close by at greater risk of health-related impacts than those living further away [15]. Also in 2012, a paper (published in this journal) documented numerous cases in which livestock and pets exposed to toxic substances from natural gas operations suffered negative health impacts and even death [16].

Public health has not been a priority for decision-makers confronting the expansion of natural gas development and consumption. Commissions to study the impacts of shale gas development have been established by Maryland and Pennsylvania and by the U.S. Secretary of Energy, but of the more than 50 members on these official bodies, none had health expertise [17]. In addition, state and federal agencies in charge of reviewing energy proposals and issuing permits do not require companies to provide information on potential health impacts, while only a few comprehensive health impact assessments (HIAs) on oil and gas development have ever been conducted in the United States [18]. Data on air and water quality near oil and gas facilities are also lacking because federal environmental testing and monitoring has long focused on a limited number of air contaminants and areas of high population density [19], while testing at oil and gas facilities in states like Pennsylvania began only recently [20]. Finally, only a few states (including Pennsylvania, Ohio, and Colorado) have any requirements for baseline air and water quality testing before drilling begins, making it difficult for researchers and regulators—as well as individuals who are directly impacted—to establish a clear connection afterwards.

SUMMARY OF THE RELEVANCE OF SELF-REPORTING HEALTH SURVEYS

For many individuals and communities living amidst oil and gas development and experiencing rapid change in their environments, too much can be at stake to rely solely on the results of long-term studies, especially those that are just now being developed. Recent examples include a new study by Guthrie Health and the Geisinger Health System in Pennsylvania, set to take from 5 to 15 years [21], and research proposals solicited in April 2012 by the National Institute of Environmental Health Sciences [22].

In contrast, self-reporting health survey research facilitates the collection and analysis of data on current exposures and medical symptoms—thereby helping to bridge the prevailing knowledge gap and pointing the way toward possible policy changes needed to protect public health. Another premise throughout the various phases of this project (location selection, survey distribution and completion, environmental testing, report development and distribution, and

58 / STEINZOR, SUBRA AND SUMI

outreach to decision-makers) was the value of public participation in science and the engagement of a variety of actors and networks to both conduct the research and ensure its beneficial application [23].

With this in mind, this health and testing project reflects some of the core principles of community-based participatory research (CBPR), including an emphasis on community engagement, use of strengths and resources within communities, application of findings to help bring about change, and belief in the research relevance and validity of community knowledge [24]. For example, the current project selected areas for investigation based in part on the observations of change in environmental conditions by long-time residents, and upon completion, participants received resources on testing and reporting of drilling problems for use in their communities.

In addition, CBPR is often used by public agencies and academic researchers to gather information on health conditions that may be related to social or environmental factors manifested on the community as well as individual level [25]. Relevant examples include identification of linkages between environmental health and socioeconomic status [26], adverse health impacts associated with coal mining [27], and the perception of health problems from industrial wind turbines [28].

Community survey and environmental testing projects such as the current one are also valuable in identifying linkages and considerations that can be used to develop protocols for additional research and policy measures. For example, community survey projects similar to the current one have revealed the presence of toxic chemicals in water and air that were known to be associated with health symptoms reported by residents, resulting in the strengthening of state standards for the control of drilling-related odors in Texas [9], expansion of a groundwater contamination investigation by the U.S. Environmental Protection Agency in Wyoming [10], and relocation of residential communities away from nearby oil refineries and contaminated waste storage areas in Louisiana [29].

METHODS

Between August 2011 and July 2012, a self-reporting health survey and environmental testing project was undertaken in order to:

- investigate the extent and types of health symptoms experienced by people living in the “gas patches” (that is, gas development areas) of Pennsylvania;
- provide air and water quality testing to some of the participating households in need of such information;
- identify possible connections between health symptoms and proximity to gas extraction and production facilities;
- provide information to researchers, officials, regulators, and residents concerned about the impact of gas development on health and air and water quality; and

- make recommendations for both further research and the development of policy measures to prevent negative health and environmental impacts.

This project did not involve certain research elements, such as structured control groups in non-impacted areas and in-depth comparative health history research, that aim to show a direct cause-and-effect relationship or to rule out additional exposures and risks. Such work, while important, was beyond the scope of the project.

The primary routes of exposure to chemicals and other harmful substances used and generated by oil and gas facilities are inhalation, ingestion, and dermal absorption—of substances in air, drinking water, or surface water—which can lead to a range of symptoms. The health survey instrument explored such variations in exposure through checklists of health symptoms grouped into categories (skin, sinus/respiratory, digestive/stomach, vision/eyes, ear/nose/mouth, neurological, urinary/urological, muscles/joints, cardiac/circulatory, reproductive, behavioral/mood/energy, lymphatic/thyroid, and immunological). A similar structure was followed for different categories of problems in participants' disease history (kidney/urological, liver, bones/joints, ulcers, thyroid/lymphatic, heart/lungs, blood disorders, brain/neurological, skin/eyes/mouth, diabetes, and cancer). Questions were also asked about occupational background and related toxic exposure history. In addition, the survey included questions on proximity to three types of facilities (compressor and pipeline stations, gas-producing wells, and impoundment or waste pits) to explore possible sources of exposure. It also asked participants to describe the type and frequency of odors they observe, since odors can both indicate the presence of a pollutant and serve as warning signs of associated health risks [30].

As indicated in Table 1, the survey was completed by 108 individuals (in 55 households) in 14 counties across Pennsylvania, with the majority (85 percent) collected in Washington, Fayette, Bedford, Bradford, and Butler counties. Taken together, the counties represent a geographical range across the state and have active wells and other facilities that have increased in number in the past few years, allowing reports of health impacts and air and water quality concerns by residents to surface [31, 32]. The survey and testing locations were all in rural and suburban residential communities.

All survey participants were assured that their names, addresses, and other identifying information on both the surveys and environmental testing results would be kept confidential and used only for purposes related to this project, such as following up with clarifying questions, responding to requests for assistance, or providing resources. Due to expressed concerns about confidentiality, participants had the option of completing the surveys anonymously, which some chose to do. Most participants answered questions on their own. In some cases, spouses, parents, or neighbors completed surveys for participants, and a few provided answers to the project coordinator in person or over the phone.

60 / STEINZOR, SUBRA AND SUMI

Table 1. Survey Locations

County surveyed	Number of surveys collected and percent of all surveys
Washington	24 (22%)
Fayette	20 (18%)
Bedford	20 (18%)
Bradford	17 (16%)
Butler	12 (11%)
Jefferson	3 (3%)
Sullivan	2 (2%)
Greene	2 (2%)
Warren	2 (2%)
Elk	2 (2%)
Clearfield	1 (1%)
Erie	1 (1%)
Susquehanna	1 (1%)
Westmoreland	1 (1%)
Total	108

While less formal and structured, the approach taken to identifying project participants has similarities to established non-random research methods that are respondent-driven and rely on word-of-mouth and a chain of referrals to reach more participants, such as “snowball” and “network” sampling [33]. As in studies in which these methods are used, the current project had a specific purpose in mind, focused on a group of people that can be hard to identify or reach, and had limited resources available for recruitment [34].

The survey was distributed in print form either by hand or through the mail and was initiated through existing contacts in the target counties. These individuals then chose to participate in the project themselves and/or recommended prospective participants, who in turn provided additional contacts. The survey was also distributed to individuals who expressed interest in participating directly to the project coordinator at public events or through neighbors, family members, and friends who had already completed surveys.

A second phase of the project involved environmental testing conducted at the homes (i.e., in the yards, on porches, or at other locations close to houses) of a

INVESTIGATING HEALTH IMPACT LINKS / 61

subset of the survey participants (70 in total) in order to identify the presence of pollutants that may be coming from gas development facilities. In all, 34 air tests and nine water tests were conducted at 35 households. Test locations were selected based on household interest, the severity of symptoms reported, and proximity to gas facilities; results were made available to the households where the testing took place. The air tests were conducted with Summa Canisters put out for 24 hours by trained individuals and the results analyzed with TO-14 and TO-15 methods, which are used and approved by the U.S. Environmental Protection Agency to test for volatile organic compounds (VOCs) such as benzene, toluene, ethylbenzene, and xylene (known as BTEX chemicals). The water tests were based on samples drawn directly from household sinks or water wells by technicians employed by certified laboratories and covered the standard Tier 1, Tier 2, and Tier 3 (including VOCs/BTEX) and in one case, gross alpha/beta radiation, radon, and radium.

FINDINGS

Health Surveys

Among participants, 45 percent were male, ranging from 18 months to 79 years of age, and 55 percent were female, ranging from 7 to 77 years of age. The closest a participant lived to gas facilities was 350 feet and the farthest away was 5 miles.

Participants had a wide range of occupational backgrounds, including animal breeding and training, beautician, child care, construction, domestic work, farming, management, mechanic, medical professional, office work, painter, retail, teaching, and welding. About 20 percent of participants reported an occupation-related chemical exposure (for example, to cleaning products, fertilizers, pesticides, or solvents). At the time of survey completion, 80 percent of participants did not smoke and 20 percent did. More than 60 percent of the current nonsmokers had never smoked, although 20 percent of nonsmokers lived with smokers.

Almost half of the survey participants answered the question on whether they had any health problems prior to shale gas development. A little less than half of those responses indicated no health conditions before the development began and a little more than half reported having had one or just a few—in particular allergies, asthma, arthritis, cancer, high blood pressure, and heart, kidney, pulmonary, and thyroid conditions were named by respondents.

While not asked specifically in the survey, some participants volunteered (verbally or in writing) additional information that points to health-related concerns warranting further investigation. For example, five reported that their existing health symptoms became worse after shale gas development started and 15 that their symptoms lessened or disappeared when they were away from home. Participants in 22 households reported that pets and/or livestock had unexplained symptoms (such as seizures or losing hair) or suddenly fell ill and died after gas development began nearby.

Some variation was noted with regard to the specific symptoms reported for each category surveyed, and some symptoms were reported to a notable degree in only one or a few locations. However, as seen in Table 2, the same overall categories of problems reported by survey participants garnered high response rates among survey participants regardless of region or county. For example, sinus/respiratory problems garnered the highest percentage of responses by participants overall, as well as in four of the five focus counties; the second top complaint category, behavioral/mood/energy, was the first in one county, second in three, and fourth in one. The total number of symptoms reported by individual participants ranged from 2 to 111; more than half reported having more than 20 symptoms and nearly one-quarter reported more than 50 symptoms. The highest numbers were reported by a 26-year-old female in Fayette County (90), a 51-year-old female in Bradford County (94), and a 59-year-old female in Warren County (111).

The 25 most prevalent individual symptoms among all participants were increased fatigue (62%), nasal irritation (61%), throat irritation (60%), sinus problems (58%), eyes burning (53%), shortness of breath (52%), joint pain (52%), feeling weak and tired (52%), severe headaches (51%), sleep disturbance (51%), lumbar pain (49%), forgetfulness (48%), muscle aches and pains (44%), difficulty breathing (41%), sleep disorders (41%), frequent irritation (39%), weakness (39%), frequent nausea (39%), skin irritation (38%), skin rashes (37%), depression (37%), memory problems (36%), severe anxiety (35%), tension (35%), and dizziness (34%).

Many symptoms were commonly reported regardless of the distance from the facility (in particular sinus problems, nasal irritation, increased fatigue, feeling weak and tired, joint pain, and shortness of breath). In addition, there was some variability in the percentage of respondents experiencing certain symptoms in relation to distance from facility, including higher rates at longer distances in a few instances. Possible influencing factors could include topography, weather conditions, participant reporting, the use of emission control technologies at facilities, or type of production (e.g., wet gas contains higher levels of liquid hydrocarbons than dry gas).

However, many symptoms showed a clearly identifiable pattern: as the distance from facilities increases, the percentage of respondents reporting the symptoms generally decreases [35]. For example, when a gas well, compressor station, and/or impoundment pit were 1500-4000 feet away, 27 percent of participants reported throat irritation; this increased to 63 percent at 501-1500 feet and to 74 percent at less than 500 feet. At the farther distance, 37 percent reported sinus problems; this increased to 53 percent at the middle distance and 70 percent at the shortest distance. Severe headaches were reported by 30 percent of respondents at the farther distance, but by about 60 percent at the middle and short distances.

Table 2. Percent of Participants Reporting Symptoms in the Most Prevalent Categories of Symptoms, by County

Symptom category	All counties	Percent of individuals reporting symptoms in category							
		Bedford	Bradford	Butler	Fayette	Washington	Others ^a		
Sinus/respiratory	88	80	82	75	85	95	87		
Behavioral/mood/energy	80	60	88	67	85	74	67		
Neurological	74	45	71	50	70	79	60		
Muscles/joints	70	55	82	67	70	74	47		
Digestive/stomach	64	55	65	58	75	63	33		
Ear/nose/mouth	66	40	59	50	75	68	47		
Skin reactions	64	45	70	67	75	63	27		
Vision/eyes	63	40	65	50	70	79	53		

^aIncludes Clearfield, Elk, Erie, Jefferson, Greene, Sullivan, Susquehanna, Warren, and Westmoreland counties. The surveys from these counties (15) were analyzed together to create a group comparable in number to each of the counties where more surveys were collected.

Figure 1 shows, for the top 20 symptoms, the percentage of residents living within 1500 feet of a natural gas facility (well, compressor, or impoundment) who reported the symptom, compared to the percentage among residents living more than 1500 feet from the facility. For 18 of the 20 symptoms, a higher percentage of those living within 1500 feet of a facility experienced the symptom than of those living farther away.

The difference in percentages reporting the symptom in the two groups (i.e., 1500 feet or closer vs. more than 1500 feet from a facility) was statistically significant for 10 of the 20 symptoms. Notably, this finding reinforces the value of data attained through self-reporting health surveys. It shows that, regardless of how symptom data were acquired, they suggest that increased proximity to gas facilities has a strong association with higher rates of symptoms reported.

When the most prevalent symptoms are broken out by age and distance from facility, some patterns stand out [35]. Within each age group, the subset living within 1500 feet of any oil and gas facility had a higher percentage of most symptoms than the age group as a whole.

Among the youngest respondents (1.5-16 years of age), for example, those within 1,500 feet experienced higher rates of throat irritation (57% vs. 69%) and severe headaches (52% vs. 69%). It is also notable that youngest group had the highest occurrence of frequent nosebleeds (perhaps reflective of the more sensitive mucosal membranes in the young), as well as experiencing conditions not typically associated with children, such as severe headaches, joint and lumbar pain, and forgetfulness.

Among 20- to 40-year-olds, those living within 1500 feet of a facility reported higher rates of nearly all symptoms; for example, 44 percent complained of frequent nosebleeds, compared to 29 percent of the entire age group. The same pattern existed among 41- to 55-year-olds with regard to several symptoms (e.g., throat and nasal irritation and increased fatigue), although with smaller differences and greater variability than in the other age groups.

The subset of participants in the oldest group (56- to 79-year-olds) living within 1,500 feet of facilities had much higher rates of several symptoms, including throat irritation (67% vs. 47%), sinus problems (72% vs. 56%), eye burning (83% vs. 56%), shortness of breath (78% vs. 64%), and skin rashes (50% vs. 33%).

In sum, while these data do not prove that living closer to oil and gas facilities causes health problems, they do suggest a strong association since symptoms are more prevalent in those living closer to facilities than those living further away. Symptoms such as headaches, nausea, and pounding of the heart are known to be the first indications of excessive exposure to air pollutants such as VOCs [36], while the higher level of nosebleeds in the youngest age group is also consistent with patterns identified in health survey projects in other states [9, 10].

The survey also asked respondents to indicate whether they were smokers. While the average number of symptoms for smokers was higher for smokers than nonsmokers (30 vs. 22), the most frequently reported symptoms were very

INVESTIGATING HEALTH IMPACT LINKS / 65




Figure 1. Association of symptoms and distance from facilities
Note: The significance of the effect was tested using a two-way contingency table analysis, and the chi-square value is given in parenthesis after each symptom. Effects significant at $p < 0.001$ are indicated by ***, those significant at $p < 0.01$ by **, and those significant at $p < 0.05$ by *.

similar (including forgetfulness, increased fatigue, lumbar pain, joint pain, eye burning, nasal irritation, sinus problems, sleep disturbances, severe headaches, throat irritation, shortness of breath, frequent nausea, muscle aches or pains, and weakness). The fact that the nonsmokers experienced symptoms that are commonly considered to be side effects of smoking (e.g., persistent hoarseness, throat irritation, sinus problems, nasal irritation, shortness of breath, and sleep disturbances) suggests that factors other than smoking were at play.

In addition, while the smoking subpopulation generally reported a larger number of symptoms, the symptoms most frequently reported by smokers and nonsmokers were remarkably similar within each age group [35]. For example, for 20- to 40-year-olds, increased fatigue, sinus problems, throat irritation, frequent nausea, and sleep problems were among the top symptoms for both smokers and nonsmokers. In the 41- to 55-year-old group, increased fatigue, throat irritation, eye burning, severe headaches, and nasal irritation were among the top symptoms for both smokers and nonsmokers, and in the over-55 age group, eye burning, sinus problems, increased fatigue, joint pain, and forgetfulness were among the top symptoms of both smokers and nonsmokers.

Participants were asked if they had noticed any odors and were asked whether they knew the source of the odors. In all but a few cases, survey participants mentioned only gas-related sources. Responses focused on locations, facilities, and processes, including drilling, gas wells, well pads, fracturing, compressor stations, condensate tanks, flaring, impoundments and pits, retention ponds, diesel engines, truck traffic, pipelines and pipeline stations, spills and leaks, subsurface ground events or migrations from underground, seismic testing, blue-colored particles in the air (possibly catalytic compounds or particulate matter), and water and stock wells. Odors were among the most common of complaints, with 81 percent of participants experiencing them sometimes or constantly. The frequency ranged from one to seven days per week and from several times per day to all day long; 18 percent said they could smell odors every day.

Participants were also asked to describe odors and whether they noticed any health symptoms when odor events occurred. The most prevalent links between odors and symptoms reported were:

- nausea: ammonia, chlorine, gas, propane, ozone, rotten gas;
- dizziness: chemical burning, chlorine, diesel, ozone, petrochemical smell, rotten/sour gas, sulfur;
- headache: chemical smell, chlorine, diesel, gasoline, ozone, petrochemical smell, propane, rotten/sour gas, sweet smell;
- eye/vision problems: chemical burning, chlorine, exhaust;
- respiratory problems: ammonia, chemical burning, chlorine, diesel, perfume smell, rotten gas, sulfur;
- nose/throat problems: chemical smell, chlorine, exhaust, gas, ozone, petrochemical smell, rotten gas, sulfur, sweet smell;

- nosebleeds: kerosene, petrochemical smell, propane, sour gas;
- skin irritation: chemical smell, chlorine, ozone, sulfur;
- decreased energy/alertness: chemical gas, ozone, rotten/sour gas, sweet smell; and
- metallic/bad taste in mouth: chemical burning, chlorine, turpentine.

Environmental Testing

As detailed in Table 3, the air tests detected a total of 19 VOCs in ambient air sampled outside of homes.

The number of compounds detected in a single sample ranged from one to 25; there was some consistency with regard to the chemicals present in most of the samples, although the concentrations of VOCs detected varied across counties [35]. The highest numbers of VOCs were detected in air samples from Washington County (15), Butler County (15), Bradford County (12), and Fayette County (9). Washington County also had the highest measured concentration of five VOCs and the second highest concentration of 12 chemicals. Samples from Butler and Bradford Counties had the highest concentrations of five and three VOCs, respectively. Five chemicals were detected in all nine of the samples from Washington County and in the six samples from Butler County: 1,1,2-trichloro-1,2,2-trifluoroethane, carbon tetrachloride, chloromethane, toluene, and trichlorofluoromethane.

It is also possible that in some places, sampling did not occur at the precise times when facilities were emitting high concentrations of chemicals or when the wind was blowing contaminants toward canisters. Some of the additional variation in number of chemicals and concentrations could be due to differences in topography, the total number of active oil and gas wells, the types of wells (conventional versus unconventional), the use of emission control technologies, and the number of active drilling sites, compressor stations, and oil and gas waste impoundments located within a certain radius of the sampling locations.

In 2010, the Pennsylvania Department of Environmental Protection (DEP) conducted air testing around natural gas wells and facilities in three regions across the state, in part using the same canister sampling methods as in this project [37]. When compared to DEP's results, our results showed some striking similarities in both the chemicals detected and concentrations. In particular, BTEX chemicals that we measured in Butler and Washington counties were consistently higher than concentrations found at DEP control sites (ethylbenzene and m- and p-xylenes were not detected at any of the control sites). When compared to the sampling done by DEP around oil and gas facilities, the concentrations in Butler and Washington counties were in the same range for benzene, but were considerably higher for toluene, ethylbenzene and m- and p-xylenes. It is also striking that some of the concentrations of ethylbenzene and

68 / STEINZOR, SUBRA AND SUMI

Table 3. Volatile Organic Compounds (VOCs) in Ambient Air, Sorted by Percent Detection^a

Compound	Total number of samples	Number of samples detecting VOCs	Percent of samples detecting VOCs	Minimum concentration	Maximum concentration	Chemical reporting limits for the three labs		
						Columbia	Con-Test	Pace ^b
2-Butanone	17	16	94	0.95	2.9	0.85-1.3	NA	NA
Acetone	17	15	88	8.0	19	6.5-10	NA	NA
Chloromethane	34	27	79	1.0	1.66	0.59-0.90	0.1	1.39-1.53
1,1,2-Trichloro-1,2,2-trifluoroethane	34	26	76	0.54	0.73	0.22-0.34	0.38	5.13-5.67
Carbon tetrachloride	34	26	76	0.4	0.76	0.091-0.14	0.31	4.21-4.65
Trichlorofluoromethane	34	26	76	0.6	1.8	0.81-1.2	0.28	3.32-3.66
Toluene	34	22	65	0.68	7.9	0.53-0.82	0.19	2.52-2.79
Dichlorodifluoromethane	17	9	63	1.9	2.8	NA	0.25	3.32-3.66
n-Hexane	8	3	38	3.03	7.04	NA	NA	2.37-2.61
Benzene	34	11	32	0.31	1.5	0.46-0.67	0.16	2.14-2.36

INVESTIGATING HEALTH IMPACT LINKS / 69

Methylene chloride	34	10	29	1.9	32.62	0.49-0.76	1.7	2.33-2.57
Total hydrocarbons (gas) ^c	8	2	25	49.8	146	NA	NA	46.9-52.2
Tetrachloroethylene	34	8	24	0.12	10.85	0.10-0.16	0.34	4.54-5.02
1,2,4-Trimethylbenzene	17	4	24	0.38	0.61	NA	0.25	3.30-3.64
Ethylbenzene	34	6	1	0.27	1.5	1.4-1.9	0.22	2.91-3.21
Trichloroethylene	34	6	18	0.17	5.37	0.08-0.12	0.27	3.60-3.98
Xylene (<i>m</i> - and <i>p</i> -)	34	5	15	0.92	5.2	2.5-3.8	0.43	2.82-3.12
Xylene (<i>o</i>)	34	5	15	0.39	1.9	1.2-1.9	0.22	2.91-3.21
1,2-Dichloroethane	34	1	3	0.64	0.64	0.59-0.90	0.2	2.71-2.99

^aConcentrations are in micrograms per cubic meter, $\mu\text{g}/\text{m}^3$ (n = total number of canister samples that were analyzed for a particular chemical).

^bPace Lab's reporting limits were in parts per billion volume (ppbv). We converted to micrograms per cubic meters ($\mu\text{g}/\text{m}^3$) using equations in the Air Unit Conversion Table (Torrent Labs, www.torrentlab.com/torrent/Home/ResourceCenter.html).

^cTotal hydrocarbons reported as parts per billion volume (ppbv).

70 / STEINZOR, SUBRA AND SUMI

xylene measured at rural and suburban residential homes in Butler and Washington counties were higher than any concentration detected by the DEP at the Marcus Hook industrial site in 2010.

As stated above, several factors can influence air results. However, it is also highly possible that the poorer air quality in the areas where we tested—which were rural and residential, with little or no other industry nearby—can be attributed to gas facilities. While the DEP reports on the agency’s air testing indicated that some of the VOCs we found in our study may not be due to oil and gas development since they persist in the atmosphere and have been widely used (for example, as refrigerants), the agency also indicates that acetone and the BTEX chemicals can be attributed to gas development [37].

With regard to the water tests conducted, Table 4 shows the 26 parameters that were detected in at least one sample. More than half of the project water samples contained methane; although some groundwater contains low concentrations of methane under normal conditions, this finding could also indicate natural gas migration from casing failure or other structural integrity problems [38]. Four of the substances detected in water well samples in Bradford and Butler Counties—manganese, iron, arsenic, and lead—were found at levels that exceed the Maximum Contaminant Levels (MCLs) set by Pennsylvania DEP’s Division of Drinking Water Management [39]. Two of the water samples, both from Butler County, were more acidic than the recommended pH for drinking water.

Some metals, such as manganese and iron, are elevated in Pennsylvania surface waters and soils, either naturally or due to past industrial activities, and levels can vary regionally [40]. In 2012, Pennsylvania State University (PSU) researchers found that some drinking water wells in the state contained somewhat elevated concentrations of certain contaminants prior to any drilling in the area [41]. However, seven out of the nine water supplies sampled in our study (78%) had manganese levels above the state MCL—a much higher percentage than what was found in the pre-drilling samples in the PSU study (27%). Even where metals are naturally occurring or predate gas development, drilling and hydraulic fracturing can contribute to elevated concentrations of these contaminants [42] and have the potential to mobilize substances in formations such as Marcellus Shale, which is enriched with barium, uranium, chromium, zinc, and other metals [43].

LINKAGES BETWEEN SURVEYS AND ENVIRONMENTAL TESTING

More research would be required to identify cause-and-effect connections between the chemicals present in air and water in Pennsylvania’s gas patches and symptoms reported by residents in specific locations. Nonetheless, such links are plausible since many of the chemicals detected in the testing are

INVESTIGATING HEALTH IMPACT LINKS / 71

known to be related both to oil and gas operations and to the health symptoms reported by individuals living at the sites where air and water testing was conducted [13-15].

The air tests together detected 19 chemicals that are known to cause sinus, skin, ear/nose/mouth, and neurological symptoms, 17 that may affect vision/eyes, and 16 that may induce behavioral effects; as well as 11 that have been associated with liver damage, nine with kidney damage, and eight with digestive/stomach problems. In addition, the brain and nervous system may be affected by five of the VOCs detected, the cardiac system by five, muscle by two, and blood cells by two [44, 45].

Using these sources [44, 45], we compared lists of the established health effects of the chemicals detected at households where testing occurred with lists of the symptoms reported in surveys by participants at those testing locations in order to identify associations. We then calculated the rate of association, in which the denominator is the total number of health impacts reported by an individual and the numerator is the total number of health impacts reported by that individual that are consistent with the known health impacts of the chemicals detected through air or water testing where they live.

Benzene, toluene, ethylbenzene, xylene, chloromethane, carbon disulfide, trichloroethylene (TCE), and acetone were detected through testing at the same households where survey participants reported symptoms established in the literature [13-15, 44, 45] as associated with these chemicals, including symptoms in the categories of sinus/respiratory, skin, vision/eyes, ear/nose/mouth, and neurological. Some of these chemicals, as well as others (such as carbon tetrachloride and tetrachloroethylene) were found at sites where survey participants reported known associated symptoms in the categories of digestion, kidney and liver damage, and muscle problems. Specific examples of chemicals and symptoms that are linked in the research literature, and were found together at households where testing and surveys were conducted, are: benzene and dizziness and nasal, eye, and throat irritation; carbon tetrachloride and nausea, headaches, and liver and kidney disease; and tetrachloroethylene and skin rashes, persistent cough, and nerve damage.

As shown in Table 5, health symptoms reported by the individuals living in a home where testing occurred matched the known health effects of chemicals detected in that home at an overall rate of 68 percent. Fayette and Washington counties had the highest match, followed by Greene, Bedford, and Butler counties.

In addition, the percent of individuals reporting symptoms that have been associated with chemicals detected in air testing at households participating in this study showed some consistency across counties with regard to the most significant categories of problems reported, as shown in Table 6—indicating that patterns in both chemicals detected and symptoms exist despite different geographic locations.

Table 4. Water Quality Results from Nine Private Water Wells in Bradford and Butler Counties, Pennsylvania

Parameter ^a	Units	Number of sample	Number above detection limit	Minimum ^b	Maximum	Mean ^c	PA DEP MCL ^d	Number of samples above MCL ^e
Barium	mg/L	9	9	0.029	0.5	0.25	2	0
Calcium	mg/L	9	9	33	66.2	43.7	None	
Magnesium	mg/L	9	9	4.5	16.8	9.1	None	
Sodium	mg/L	9	9	9.2	64.1	20.9	None	
Strontium	mg/L	9	9	0.126	1.7	0.5	None	
Hardness (total as CaCO ₃)	mg/L	9	9	120	234	147	None	
pH	Std Units	9	9	6	7.9	6.5	6.5-8.5	<i>f</i>
Alkalinity (total as CaCO ₃)	mg/L	9	9	38	285	130	None	
Total dissolved solids	mg/L	9	9	138	392	218	500	0
Sulfate	mg/L	9	9	6.7	231	33	250	0
Manganese	mg/L	9	7	< 0.005	6.44	1.04	0.05	7
Chloride	mg/L	9	7	< 5.0	84.3	24.1	250	0
Iron	mg/L	9	6	< 0.04	153	19.5	0.3	5
Potassium	mg/L	6	6	1.14	1.57	1.1	None	

INVESTIGATING HEALTH IMPACT LINKS / 73

Specific conductance	µmhos/cm	6	6	287	552	326	None
Methane	µg/L	9	5	1.06	57.4	10	0.3
Arsenic	mg/L	9	4	< 0.001	0.0282	0.005	0.010
Lead	mg/L	9	4	< 0.001	0.113	0.113	0.01
Total coliform	per 100 mL	9	4	Absent	Present		None
Total suspended solids	mg/L	6	4	< 5	448	118	None
Temperature, water	Degree/Celsius	3	3	25	29	28	None
Turbidity	NTU	3	3	0.22	5.7	2.3	None
Nitrate	mg/L	3	3	0.076	0.71	0.46	10
<i>E. coli</i>	per 100 mL	9	2	Absent	Present		None
Sulfur	µg/L	1	1	< 1,000	7,550	2,850	None
Bromide	mg/L	1	1	0.26	0.26	0.26	None

^aNote: not all parameters were analyzed in every sample.

^bMinimum values: If reports included non-detects of a particular chemical, the minimum value in the table was shown as being less than (<) the lowest laboratory detection limit.

^cMean values: Non-detected chemicals were assigned a concentration equal to half of the detection limit *only if* there were other samples that detected the chemical.

^dMCL: Maximum Contaminant Levels published by the Pennsylvania Department of Environmental Protection Division of Drinking Water Management.

^eNo values are provided if MCLs for substances do not exist.

^fTwo samples had higher acidity (lower pH) than the value recommended by the PA DEP.

74 / STEINZOR, SUBRA AND SUMI

Table 5. Match between Health Symptoms Reported by Individuals at Air Testing Sites and Known Effects of Chemicals Detected

County	Number of individuals surveyed at homes where testing was conducted	Match between known health effects of chemicals detected and symptoms reported (percent) ^a	
		Average	Range
Overall	59	68	33-100
Fayette	16	73	33-100
Washington	15	73	33-100
Bradford	8	58	16-100
Butler	8	63	56-68
Bedford	6	69	63-100
Elk	2	64	53-74
Clearfield	1	none	none
Greene	1	70	70
Susquehanna	1	50	50

^aWhen a health symptom was associated in the literature with more than one of the chemicals detected, only one match was counted for that symptom.

As mentioned above, levels of iron, manganese, arsenic, and lead were detected in our water well samples in Bradford and Butler Counties at levels that exceeded drinking water standards set by the Pennsylvania DEP. These substances are known to be associated with numerous symptoms reported by individuals living in the homes where these particular exceedances occurred, including symptoms in the categories of sinus/respiratory, skin reactions, digestive/stomach, vision/eyes, ear/nose/mouth, neurological, muscle/joint, behavioral/mood/energy, and liver and kidney damage. Survey participants in the homes where water samples contained methane reported health symptoms known to be associated with methane, including symptoms in the categories of sinus/respiratory, digestive/stomach, neurological, and behavioral/mood/energy. While the water samples taken for this project did not show detectable exceedances of safety standards for other substances, it is notable that no drinking water standards have been set for methane, bromide, sodium, strontium, or Total Suspended Solids (TSS)—and thus no exceedances would be indicated in laboratory reports.

Table 6. Percent of Individuals at Air Testing Sites Reporting Symptoms Associated in the Literature with Chemicals Detected at Those Sites, by Symptom Category and Primary Air

Symptom category	Testing counties							
	All	Bedford	Bradford	Butler	Fayette	Washington	Others ^a	
Sinus/respiratory	83	100	88	100	81	73	80	
Vision/eyes	73	—	100	63	69	67	60	
Digestive/stomach	69	50	63	88	75	80	—	
Skin reactions	63	50	63	88	69	53	40	
Neurological	60	50	88	75	44	53	60	
Behavioral/mood/energy	54	67	50	63	63	47	40	
Ear/nose/mouth	33	50	—	38	44	33	20	
Muscle problems	—	—	—	—	—	40	—	

^aThis includes air samples from Clearfield, Elk, Greene, and Susquehanna counties.

DISCUSSION

Complete evidence regarding health impacts of gas drilling cannot be obtained due to incomplete testing and disclosure of chemicals, and non-disclosure agreements. Without rigorous scientific studies, the gas drilling boom sweeping the world will remain an uncontrolled health experiment on an enormous scale.

—Michelle Bamberger and Robert Oswald [16]

While the survey and testing results, and their related findings, do not constitute definitive proof of cause and effect, we believe they do indicate the strong likelihood that the health of people living in proximity to gas facilities is being affected by exposure to pollutants from those facilities. Most participants report a high number of health symptoms; similar patterns of symptoms were identified across project locations and distances from facilities; and consistency in symptoms reported exists regardless of age group or smoking history. In addition, contaminants that result from oil and gas development were detected in air and water samples in areas where residents are experiencing health symptoms that are established in the literature as consistent with such exposures.

Because of the short-term nature of the air-canister testing (24 hours) and the single water tests conducted at households, our results were contingent on conditions at particular “moments in time.” Thus additional chemicals, or the same chemicals at different concentrations, might be captured through expanded testing; and residents could be experiencing exposures that were not detected but would be detectable through such testing. In addition, some of the variation in the air test results may have been due to the different reporting protocols used by the laboratories used in this project. Although all the labs test for the same core suite of chemicals, both their reporting limits and the additional chemicals for which they test vary; these will be key considerations for future testing work.

Another consideration that warrants further exploration involves the established standards on both the state and federal levels for “safe” concentrations, which are set only for exposure to single contaminants. This prevailing regulatory approach can not adequately address the potential risks posed by chronic, long-term exposure to lower levels of multiple contaminants simultaneously—in other words, the experience of people living in oil and gas areas day in and day out, and of workers at job sites where toxic substances are continuously used. In addition, for many substances in the environment (including those that come from gas operations and were detected in our air and water sampling), data on health risks or safe exposure levels simply do not exist.

More research is also needed that focuses on the sources of odors and odor events experienced by residents living near gas facilities. In some cases, participants reported different health impacts associated with specific sources and odor events than those they reported in the overall health survey. Since odors are

INVESTIGATING HEALTH IMPACT LINKS / 77

a clear sign of the presence of airborne substances (such as fuel and chemicals), this aspect warrants tracking and analysis.

Although we did not investigate additional factors that can influence health conditions (e.g., through ordered control groups, in-depth health history research, or identification of other potential sources of contaminants), such factors may affect an individual's health independent of gas operations. The relationship between symptoms and distance from gas facilities also warrants more research.

At the same time, we strongly suggest that for individuals with a history of other health concerns (e.g., asthma or heart conditions) and who are already living with other exposures (e.g., traffic fumes or workplace chemicals), the presence of gas facilities and related pollution could have a strong “trigger effect” that can make existing problems worse and put individuals at higher risk of developing new ones.

RECOMMENDATIONS

As discussed earlier, scientific knowledge about the health and environmental impacts of shale gas development—and also the adoption of policy and regulatory measures to prevent them—are proceeding at a far slower pace than the development itself. This timing mismatch creates situations (already being experienced by residents of Pennsylvania and other states) in which problems are widely reported but left unaddressed. Several measures can be taken to ensure that public health impacts are fully understood and given greater priority in decision-making about shale gas development.

1) Elevate the role of public health considerations in gas development decisions. A key measure would be to conduct health impact assessments before permitting begins. HIAs aim to minimize negative impacts and to improve health outcomes associated with land use decisions by analyzing problems that could arise over time as well as existing health and environmental risks that could be exacerbated by new activities [46]. HIAs can also have a strong preventive effect by identifying mitigation measures related to aspects such as toxic exposures, air and water pollution, and emergency response [47]. In addition, regulatory agencies could comprehensively plan the scope and pace of permits for wells and other facilities in order to reduce impacts on air and water quality, rather than continuing the permit-by-permit process currently being followed in Pennsylvania and other states. Information on where wells and facilities would be built in relation to places where health could be at risk (e.g., homes, schools, and hospitals) could also be required in permit applications.

2) Increase the involvement of state departments of health in assessing the impacts of gas development. Efforts should be increased to track and respond to health concerns, and a database should be established to document these problems and the agency response. Health departments could provide training for health and medical professionals on exposure pathways and health symptoms

78 / STEINZOR, SUBRA AND SUMI

related to gas operations, so that residents receive more informed advice and appropriate testing and care referrals. Financial aid mechanisms should be established to enable low-income residents to have blood and urine tests for chemical exposure.

3) Conduct baseline water testing and continuous long-term monitoring of air quality. Such testing would apply to private wells and public drinking water supplies prior to drilling and to the air at or near facilities during all phases of operations. Testing and monitoring should cover a full suite of chemicals, and contaminants and results should be reported regularly and made available to the public. Air quality testing in particular should be conducted at a range of facilities (e.g., compressor stations, impoundment pits, dehydrators) that cause emissions. These efforts could be carried out by the state regulatory agencies that issue permits or through an agreement between those agencies and health departments. Inter-agency agreements could also be developed to track potential health impacts that could result following spills of chemicals and waste, the underground migration of fracturing fluids, leaks, and other problems.

4) Strengthen regulations for facilities to minimize air and water pollution risks. These could include significantly increased setback distances; the installation of advanced technologies on all equipment to reduce emissions, odors, and noise; the use of closed-loop storage systems for waste and drilling fluids (rather than open pits); and the practice of “green completions” to reduce or eliminate flaring and venting of methane gas and other pollutants.

5) Advance changes in testing parameters that determine safe exposure in order to account for low-level, chronic exposure and multiple chemical exposure in testing and monitoring. Such changes are necessary to reflect impacts on people living in oil and gas development areas day in and day out, as well as workers at facilities. Under current testing parameters (which are based largely on acute episodes involving single contaminants), results may show below-threshold levels even though residents are negatively affected. For example, a recent paper showed that endocrine-disrupting chemicals can have different but still harmful effects at lower doses than at higher ones and concluded that fundamental changes in chemical testing and safety protocols are needed to protect human health [48]. Additionally, current health guidelines should be updated to capture more of the chemicals currently in use and to assess complex or indirect sources of contamination, such as oil and gas operations that rely on a variety of substances, equipment, and facilities at numerous stages of development.

CONCLUSION

While we realize that human activities may involve hazards, people must proceed more carefully than has been the case in recent history. Corporations, government entities, organizations, communities, scientists, and other individuals must adopt a precautionary approach to all human endeavors. . . . When an activity raises threats of harm to human health or the environment,

INVESTIGATING HEALTH IMPACT LINKS / 79

precautionary measures should be taken even if some cause and effect relationships are not fully established scientifically.

—Wingspread Consensus Statement on the Precautionary Principle [49]

Across the gas patches of the United States, people experiencing health problems voice the simple wish to be believed. Many say that their health has worsened since gas development began in their communities and that they feel better when they are away from home. Often these conversations turn to what it will take for regulators and policymakers to view their stories not just as “anecdotes,” but as valid concerns worthy of an effective response.

There is no doubt that more research on the environmental and health impacts of shale gas development is needed and can play a critical role in making sound decisions about a complex and controversial issue. Yet an equally important consideration is how to respond to the presence of unanswered questions. For many proponents of unfettered gas development, the absence of definitive causal links between gas facilities and specific health impacts indicates the absence of a problem. But for impacted communities and others who believe health and the environment deserve protection and that water and air quality should be maintained, what we don’t yet know makes the need for caution even greater.

We believe that the findings of this survey and testing project in Pennsylvania, coupled with similar projects elsewhere and an emerging body of research, provide sufficient evidence for decision-makers to take action to slow the rush to drill, at least until the wide gaps in scientific knowledge, policies, and regulations are bridged. Much is already known about the chemicals used and pollution caused by oil and gas activities, which alone create the real potential for negative health effects in any area where development occurs [50]. The precautionary principle should be applied to decisions about shale gas development (both in existing gas patches and in areas slated for new development), and this should include shifting the burden of proof that harm does or does not occur to those proposing the action.

The status quo—in which science and policy changes proceed slowly while gas development accelerates rapidly—is likely to worsen air and water quality, resulting in negative health impacts and possibly a public health crisis. Greater understanding of the experiences reported by individuals living near gas facilities can play an important role in pointing the way forward to preventing these problems, both in Pennsylvania and nationwide.

ACKNOWLEDGMENTS

This project would not have been possible without the caring, concern, and openness of project participants, who shared their time and personal experiences and trusted us to write about them. Much gratitude is also owed to the community members and organizational partners who provided the contacts and guidance

80 / STEINZOR, SUBRA AND SUMI

that made it possible to engage project participants. Many thanks to the Colcom Foundation for its generous support of this project and commitment to protecting the environment and public health.

AUTHORS' BIOGRAPHIES

NADIA STEINZOR is the Eastern Program Coordinator of Earthworks' Oil & Gas Accountability Project. She works with landowners, concerned citizens, and environmental organizations to educate and support impacted communities, advance state oil and gas regulations, and promote Earthworks' federal reform agenda. Nadia has worked for nearly two decades in research, communications, writing, and editing to promote action and policies on environmental protection and land conservation. She holds an M.S. in Environmental Policy from Bard College and an M.A. in Peace and Development Studies from the University of Gothenburg, Sweden. Send email to her at nsteinzor@earthworksaction.org

WILMA SUBRA is President of Subra Company, Inc., a chemistry lab and environmental consulting firm in New Iberia, Louisiana, which provides technical assistance to citizens and victims of environmental issues. She serves on local, state, and federal committees dealing with oilfield waste, oil refineries, and toxic release inventory. Wilma holds an M.S. in Microbiology and Chemistry from the University of Southwestern Louisiana. She is a recipient of the MacArthur Fellowship Genius Award for helping ordinary citizens understand, cope with, and combat environmental problems in their communities. Send email to her at subracom@aol.com

LISA SUMI is a consultant who works on the environmental, community, and health impacts of extractive industries such as coal and hard rock mining, tar sands, and oil and gas. She conducts work that integrates science, technology, policy, and corporate research and analysis with community education and outreach. She has served as Research Director for the Oil and Gas Accountability Project and the Environmental Mining Council of British Columbia, and has worked for clients such as As You Sow, Council of Canadians, Earthworks, ForestEthics, Initiative for Responsible Mining Assurance, Massachusetts Institute of Technology Center for Future Civic Media, MiningWatch Canada, and Natural Resources Defense Council. Lisa holds a M.S. in Physical Geography from the University of Toronto. Isumi@earthworksaction.org

NOTES

1. Institute of Medicine, "The Health Impact Assessment of New Energy Sources: Shale Gas Extraction," www.iom.edu/Activities/Environment/EnvironmentalHealthRT/2012-APR-30.aspx (accessed June 27, 2012).

INVESTIGATING HEALTH IMPACT LINKS / 81

2. U.S. Energy Information Administration, "Number of Producing Gas Wells," [data table], http://205.254.135.7/dnav/ng/ng_prod_wells_s1_a.htm (accessed July 8, 2012).
3. Pennsylvania Department of Environmental Protection, "Oil and Gas Reports," www.portal.state.pa.us/portal/server.pt/community/oil_and_gas_reports/20297#InteractiveReports (accessed September 10, 2012).
4. C. W. Schmidt, "Blind Rush? Shale Gas Boom Proceeds amid Human Health Questions," *Environmental Health Perspectives* 119 (2011): A348-353, doi: 10.1289/ehp.119-a348.
5. Earthworks Oil and Gas Accountability Project, "Oil and Gas Pollution," www.earthworksaction.org/issues/detail/oil_and_gas_pollution (accessed September 20, 2012).
6. Pennsylvania Alliance for Clean Water and Air, "List of the Harmed," <http://pennsylvaniaallianceforcleanwaterandair.wordpress.com/the-list/> (accessed July 8, 2012).
7. Edward Humes, "Fractured Lives: Detritus of Pennsylvania's Shale Gas Boom," *Sierra Magazine*, July/August 2012, 52-59.
8. Earthworks' Oil & Gas Accountability Project, *Natural Gas Flowback: How the Texas Gas Boom Affects Community Health and Safety*, 2011.
9. Earthworks' Oil & Gas Accountability Project, *Health Survey Results of Current and Former Residents of DISH/Clark, Texas Residents*, 2009.
10. Earthworks' Oil & Gas Accountability Project, *Community Health Survey Results: Pavillion, Wyoming*, 2010.
11. Subra Company, *Health Report on Mossville, Calcasieu Parish, Louisiana*, 2009.
12. Louisiana Environmental Action Network, "Survey of the Human Health Impacts Due to the BP Deepwater Horizon Disaster," 2012, <http://leanweb.org/our-work/water/bp-oil-spill/results-of-the-louisiana-environmental-action-network-lean-survey-of-the-human-health-impacts-due-to-the-bp-deepwater-horizon-disaster> (accessed July 15, 2012).
13. R. Witter et al., *Potential Exposure-Related Human Health Effects of Oil and Gas Development: A Literature Review (2003-2008)*, 2008.
14. T. Colborn et al., "Natural Gas Operations from a Public Health Perspective," *Human & Ecological Risk Assessment* 17 (5) (2011): 1039-1056, doi: 10.1080/10807039.2011.605662.
15. L. M. McKenzie et al., "Human Health Risk Assessment of Air Emissions from Development of Unconventional Natural Gas Resources," *Science of the Total Environment* 1 (424) (2012): 79-87, doi: 10.1016/j.scitotenv.2012.02.018.
16. M. Bamberger and R. E. Oswald, "Impacts of Gas Drilling on Human and Animal Health," *New Solutions: A Journal of Environmental and Occupational Health Policy* 22 (1) (2012): 51-77, doi: 10.2190/NS.22.1.e.
17. B. Goldstein, J. Kriesky, and B. Pavliakova, "Missing from the Table: Role of the Environmental Public Health Community in Governmental Advisory Commissions Related to Marcellus Shale Drilling," *Environmental Health Perspectives* 120 (4) (2012): 483-486, doi: 10.1289/ehp.1104594.
18. The Health Impact Project, "HIA in the United States," www.healthimpactproject.org/hia/us (accessed June 27, 2012).
19. U.S. Environmental Protection Agency, "Ambient Air Monitoring Program," <http://www.epa.gov/airquality/qa/monprog.html> (accessed September 20, 2012).

82 / STEINZOR, SUBRA AND SUMI

20. Clean Air Council, "PA DEP Launches Long-term Air Sampling," http://cleanair.org/program/outdoor_air_pollution/marcellus_shale/pa_dep_launches_long_term_air_sampling (accessed September 20, 2012).
21. Geisinger Health System, "Guthrie Health and Geisinger Collaborate on Marcellus Shale Work," <https://webapps.geisinger.org/ghsnews/articles/GuthrieHealthandGeisingerc8464.html> (accessed September 22, 2012).
22. National Institutes of Environmental Health Sciences, "Shale Gas Workshop Explores Needed Research on Fracking," *Environmental Factor*, June 2012, www.niehs.nih.gov/news/newsletter/2012/6/spotlight-shalegas/index.htm (accessed September 22, 2012).
23. B. L. Allen, "Shifting Boundary Work: Issues and Tensions in Environmental and Health Science in the Case of Grand Bois, Louisiana," *Science as Culture* 13 (12) (2004): 429-448, doi: 10.1080/0950543042000311805.
24. Minkler, M. et al., *Community-Based Participatory Research: A Strategy for Building Health Communities and Promoting Health through Policy Change*, PolicyLink, School of Public Health, University of California Berkeley, 2012.
25. B. A. Israel et al., "Review of Community-Based Research: Assessing Partnership Approaches to Improve Public Health," *Annual Review of Public Health* 19 (1998): 173-202, doi: 10.1146/annurev.publhealth.19.1.173.
26. National Institute of Environmental Health Sciences, "Community-Based Participatory Research in Environmental Health," Request for Applications ES-01-003, January 2001.
27. M. Hendryx et al., "Self-Reported Cancer Rates in Two Rural Areas of West Virginia With and Without Mountaintop Coal Mining," *Journal of Community Health* 37 (4) (2012): 32-327, doi: 10.1007/s10900-011-9448-5.
28. C. M. E. Krogh et al., "WindVOiCe, A Self-Reporting Survey: Adverse Health Effects, Industrial Wind Turbines, and the Need for Vigilance Monitoring," *Bulletin of Science Technology Society* 31 (4) (2011): 334-345, doi: 10.1177/0270467611412551.
29. Mossville Environmental Action Now, Subra, W., and Advocates for Environmental Human Rights, *Industrial Sources of Dioxin Poisoning in Mossville, Louisiana: A Report Based on the Government's Own Data*, 2007.
30. S. Schiffman, "Science of Odors as a Potential Health Issue," *Journal of Environmental Quality* 34 (2005): 129-138, doi: 10.2134/jeq2005.0129.
31. Pennsylvania Department of Environmental Protection, eFacts database, www.ahs2.dep.state.pa.us/eFACTSWeb/criteria_sitesbymuni.aspx (accessed October 3, 2012).
32. FracTracker, Data-Generated Map of the Distribution of Gas Wells across Pennsylvania, www.fractracker.org/maps (accessed October 3, 2012).
33. D. D. Heckathorn, "Respondent-Driven Sampling: A New Approach to the Study of Hidden Populations," *Social Problems* 44 (2c) (1997): 174-199.
34. Department of Psychology, University of California at Davis, "Types of samples," <http://psychology.ucdavis.edu/sommerb/sommerdemo/sampling/types.htm> (accessed June 25, 2012).
35. Earthworks' Oil & Gas Accountability Project, *Gas Patch Roulette: How Shale Gas Development Risks Public Health in Pennsylvania*, 2012.
36. Swiss Agency for the Environment, Forests, and Landscapes, "National Air Pollutions Monitoring Network," 2001.

INVESTIGATING HEALTH IMPACT LINKS / 83

37. Pennsylvania Department of Environmental Protection, Northeastern Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report; Northcentral Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report, 2011; and Southwestern Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report, 2010.
38. A. E. Ingraffea, Fluid Migration Mechanisms Due to Faulty Well Design and/or Construction: An Overview and Recent Experiences in the Pennsylvania Marcellus Play, 2012.
39. Pennsylvania Department of Environmental Protection, Division of Drinking Water Management, "Maximum Contaminant Levels," www.portal.state.pa.us/portal/server.pt?open=18&objID=437830&mode=2 (accessed July 20, 2012).
40. National Science Foundation, "Can Marcellus Shale Development and Healthy Waterways Co-exist?," www.nsf.gov/discoveries/disc_summ.jsp?cntn_id=122543 (accessed October 2, 2012).
41. Pennsylvania State University Extension. "Iron and Manganese in Private Water Systems," <http://extension.psu.edu/water/converted-publications/iron-and-manganese-in-private-water-systems> (accessed October 2, 2012).
42. E. Boyer et al., The Center for Rural Pennsylvania, Pennsylvania State University, The Impact of Marcellus Gas Drilling on Rural Drinking Water Supplies, 2011.
43. T. Banks, "Fluid Rock Interactions Associated with Hydraulic Fracturing and Natural Gas Development," Global Water Program Magazine, January 2011, http://globalwater.jhu.edu/index.php/magazine/article/fluid_rock_interactions_associated_with_hydraulic_fracturing_and_natural_gas/ (accessed October 2, 2012).
44. Agency for Toxic Substances and Disease Registry (ATSDR), "Toxicological Profiles," <http://www.atsdr.cdc.gov/toxprofiles/index.asp> (accessed July 30, 2012).
45. New Jersey Department of Health, "Right to Know Hazardous Substance Fact Sheets," <http://web.doh.state.nj.us/rtkhsfs/indexfs.aspx> (accessed July 30, 2012).
46. A. Wernham, "Health Impact Assessment for Shale Gas Extraction," www.healthimpactproject.org/resources/health-impact-assessment-for-shale-gas-extraction (accessed July 10, 2012).
47. Garfield County, Colorado, "Battlement Mesa Health Impact Assessment," www.garfield-county.com/environmental-health/battlement-mesa-health-impact-assessment-ehms.aspx (accessed July 10, 2012).
48. L. N. Vandenberg et al., "Hormones and Endocrine-Disrupting Chemicals: Low-Dose Effects and Nonmonotonic Dose Responses," *Endocrine Reviews* 33 (3) (2012): 1-78, doi: 10.1210/er.2011-1050.
49. Science and Environmental Health Network, "Wingspread Consensus Statement on the Precautionary Principle," 1998, www.sehn.org/wing.html (accessed July 18, 2012).
50. R. Witter et al., Potential Exposure-Related Human Health Effects of Oil and Gas Development: A White Paper, 2008.

Direct reprint requests to:

Nadia Steinzor
Earthworks' Oil & Gas Accountability Project
P.O. Box 149
Willow, NY 12495
e-mail: nsteinzor@earthworksaction.org